

Tyagi LK, Kori ML. Formulation and *in vitro* evaluation of Eudragit® RS 100 microspheres containing lornoxicam prepared by emulsion-solvent evaporation method. *Bull. Pharm. Res.* 2013;3(3):112-20.

**References (30):**

1. Alex R, Bodmeier R. Encapsulation of water-soluble drugs by a modified solvent evaporation method. I. Effect of process and formulation variables on drug entrapment. *J. Microencapsul.* 1990;7(3):347-55.  
<http://www.ncbi.nlm.nih.gov/pubmed/2384837>
2. Atyabi F, Mohammadi A, Dinarvand R. Preparation of nimodipine loaded microspheres: Evaluation of parameters. *Iranian J. Pharm. Sci.* 2005;1(3):143-52.  
[http://www.sid.ir/en/VIEWSSID/J\\_pdf/106720050303.pdf](http://www.sid.ir/en/VIEWSSID/J_pdf/106720050303.pdf)
3. Babay D, Hoffman A, Benita S. Design and release kinetic pattern evaluation of indomethacin microspheres intended for oral administration. *Biomaterials* 1988;9(6): 482-8.  
<http://www.ncbi.nlm.nih.gov/pubmed/3224134>
4. Balfour JA, Fitton A, Barradell LB. Lornoxicam. A review of its pharmacology and therapeutic potential in the management of painful and inflammatory conditions. *Drugs* 1996;51(4):639-57.  
<http://www.ncbi.nlm.nih.gov/pubmed/8706598>
5. Basarkar GD, Shirsath GN, Patil SB. Development of microspheres containing diclofenac diethylamine as sustained release topical formulation. *Bull. Pharm. Res.* 2013;3(1):14-22.  
[www.appconnect.in/wp-content/uploads/2011/03/ReprintBPR0662.pdf](http://www.appconnect.in/wp-content/uploads/2011/03/ReprintBPR0662.pdf)
6. Basu SK, Adhiyaman R. Preparation and characterization of nitrendipine loaded eudragit RL 100 microspheres prepared by an emulsion-solvent evaporation method. *Trop. J. Pharm. Res.* 2008;7(3):1033-41.  
[http://www.tjpr.org/vol7\\_no3/735\\_rajendran.php](http://www.tjpr.org/vol7_no3/735_rajendran.php)
7. Behera BC, Sahoo SK, Dhal S, Barik BB, Gupta BK. Characterization of glipizide-loaded polymethacrylate microspheres prepared by an emulsion solvent evaporation method. *Trop. J. Pharm. Res.* 2008;7(1):879-85.  
[http://www.tjpr.org/vol7\\_no1/713Behera.php](http://www.tjpr.org/vol7_no1/713Behera.php)
8. Choi HS, Seo SA, Khang G, Rhee JM, Lee HB. Preparation and characterization of fentanyl-loaded PLGA microspheres: *In vitro* release profiles. *Int. J. Pharm.* 2002;234(1-2):195-203.  
<http://www.ncbi.nlm.nih.gov/pubmed/11839450>

9. Chowdary KPR, Koteswara Rao N, Malathi K. Ethyl cellulose microspheres of glipizide: Characterization, *in vitro* and *in vivo* evaluation. *Indian J. Pharm. Sci.* 2004;66(4):412-6.  
<http://www.ipsonline.com/article.asp?issn=0250-474X;year=2004;volume=66;issue=4;spage=412;epage=416;aulast=Chowdary;type=0>
10. Dahiya S, Gupta ON. Formulation and *in vitro* evaluation of metoprolol tartrate microspheres. *Bull. Pharm. Res.* 2011;1(1):31-9.  
[www.appconnect.in/wp-content/uploads/2012/01/ReprintBPR007.pdf](http://www.appconnect.in/wp-content/uploads/2012/01/ReprintBPR007.pdf)
11. Garud N, Garud A. Preparation and *in-vitro* evaluation of metformin microspheres using non-aqueous solvent evaporation technique. *Trop. J. Pharm. Res.* 2012;11(4):577-83.  
<http://www.aiol.info/index.php/tjpr/article/view/80566>
12. Gupta VK, Beckert TE, Price JC. A novel pH- and time-based multi-unit potential colonic drug delivery system. I. Development. *Int. J. Pharm.* 2001;213(1-2):83-91.  
<http://www.ncbi.nlm.nih.gov/pubmed/11165096>
13. Haznedar S, Dortunc B. Preparation and *in vitro* evaluation of eudragit microspheres containing acetazolamide. *Int. J. Pharm.* 2004;269(1):131-40.  
<http://www.ncbi.nlm.nih.gov/pubmed/14698584>
14. Homdrum E-M, Likar R, Nell G. Xefo® Rapid: A novel effective tool for pain treatment. *Eur. Surg.* 2006;38(5):342-52.  
<http://link.springer.com/article/10.1007%2Fs10353-006-0272-6#>
15. Khan SA, Ahmad M, Murtaza G, Aamir MN, Rehman N, Kousar R, Rasool F, Akhtar M. Formulation of nimesulide floating microparticles using low-viscosity hydroxypropyl methylcellulose. *Trop. J. Pharm. Res.* 2010;9(3):293-9.  
[http://www.tjpr.org/vol9\\_no3/2010\\_9\\_3\\_11.php](http://www.tjpr.org/vol9_no3/2010_9_3_11.php)
16. Kidd B, Frenzel W. A multicenter, randomized, double blind study comparing lornoxicam with diclofenac in osteoarthritis. *J. Rheumatol.* 1996;23(9):1605-11.  
<http://www.ncbi.nlm.nih.gov/pubmed/8877932>
17. Kim BK, Hwang SJ, Park JB, Park HJ. Preparation and characterization of drug-loaded polymethacrylate microspheres by an emulsion solvent evaporation method. *J. Microencapsul.* 2002;19(6):811-22.  
<http://www.ncbi.nlm.nih.gov/pubmed/12569029>
18. Kumar M, Dureja H. Development and characterization of factorially designed 5-fluoro uracil microspheres. *Bull. Pharm. Res.* 2011;1(1):54-61.  
[www.appconnect.in/wp-content/uploads/2012/01/FPP-BPR010.pdf](http://www.appconnect.in/wp-content/uploads/2012/01/FPP-BPR010.pdf)
19. Maia JL, Santana MHA, Re MI. The effect of some processing conditions on the characteristics of biodegradable microspheres obtained by an emulsion solvent evaporation process. *Braz. J. Chem. Eng.* 2004;21(1):1-12.  
[http://www.scielo.br/scielo.php?pid=S0104-66322004000100002&script=sci\\_arttext](http://www.scielo.br/scielo.php?pid=S0104-66322004000100002&script=sci_arttext)

20. Martin AN. Physical Pharmacy: Physical Chemical Principles in the Pharmaceutical Sciences, 4<sup>th</sup> edition, Lippincott Williams & Wilkins: USA, 1993.
21. Perumal D. Microencapsulation ibuprofen and eudragit® RS 100 by the emulsion solvent diffusion technique. *Int. J. Pharm.* 2001;218(1-2):1-11.  
<http://www.ncbi.nlm.nih.gov/pubmed/11337145>
22. Pongpaibul Y, Price JC, Whitworth CW. Preparation and evaluation of controlled release indomethacin microspheres. *Drug Dev. Ind. Pharm.* 1984;10(10):1597-1616.  
<http://informahealthcare.com/doi/abs/10.3109/03639048409039070>
23. Ritschel WA. Biopharmaceutics and pharmacokinetic aspects in the design of controlled release peroral drug delivery systems. *Drug Dev. Ind. Pharm.* 1989;15(6-7):1073-1103.  
<http://informahealthcare.com/doi/abs/10.3109/03639048909043666?journalCode=ddi>
24. Sahoo SK, Mallick AA, Barik BB, Senapati PC. Formulation and *in vitro* evaluation of eudragit® microspheres of stavudine. *Trop. J. Pharm. Res.* 2005;4(1):369-75.  
<http://www.ajol.info/index.php/tjpr/article/view/14622>
25. Sahoo SK, Mallick AA, Barik BB, Senapati PC. Preparation and *in vitro* evaluation of ethyl cellulose microspheres containing stavudine by the double emulsion method. *Pharmazie* 2007;62(2):117-21.  
<http://www.ncbi.nlm.nih.gov/pubmed/17341031>
26. Sengel CT, Hascicek C, Gonul N. Development and *in-vitro* evaluation of modified release tablets including ethylcellulose microspheres loaded with diltiazem hydrochloride. *J. Microencapsul.* 2006;23(2):135-52.  
<http://www.ncbi.nlm.nih.gov/pubmed/16754371>
27. Singh V, Chaudhary AK. Preparation of Eudragit E100 microspheres by modified solvent evaporation method. *Acta Pol. Pharm.* 2011;68(6):975-80.  
<http://www.ncbi.nlm.nih.gov/pubmed/22125964>
28. Song SZ, Cardinal JR, Kim SM, Kim SW. Progestin permeation through polymer membrane. V: Progesterone release from monolithic hydrogel devices. *J. Pharm Sci.* 1981;70(2):216-9.  
<http://www.ncbi.nlm.nih.gov/pubmed/7205230>
29. Skjeldt NM, Davies NM. Clinical pharmacokinetics of lornoxicam. A short half-life oxicam. *Clin. Pharmacokinet.* 1998;34(6):421-8.  
<http://www.ncbi.nlm.nih.gov/pubmed/9646006>
30. Takka S, Rajbhandari S, Sakr A. Effect of anionic polymers on the release of propranolol hydrochloride from matrix tablets. *Eur. J. Pharm. Biopharm.* 2001;52(1):75-82.  
<http://www.ncbi.nlm.nih.gov/pubmed/11438426>