

Dahiya S, Tayde P. Binary and ternary solid systems of carvedilol with 2-hydroxypropyl- $\beta$ -cyclodextrin and PVP K30. *Bull. Pharm. Res.* 2013;3(3):128-34.

### References (14):

1. Bekers O, Uijtendal EV, Beijnen JH, Bult A, Underberg WJ. Cyclodextrins in pharmaceutical field. *Drug Dev. Ind. Pharm.* 1991;17(11):1503-49.  
<http://informahealthcare.com/doi/abs/10.3109/03639049109026630>
2. Dahiya S, Pathak K. Dissolution enhancement of aceclofenac by  $\beta$ -cyclodextrin complexation. *J. Pharm. Res.* 2006a;5(4):99-103.
3. Dahiya S, Pathak K. Physicochemical characterization and dissolution enhancement of aceclofenac-hydroxypropyl  $\beta$ -cyclodextrin binary systems. *PDA J. Pharm. Sci. Tech.* 2006b; 60(6):378-88.  
<http://journal.pda.org/content/60/6/378.abstract>
4. Dahiya S, Pathak K. Influence of amorphous cyclodextrin derivatives on aceclofenac release from directly compressible tablets. *Pharmazie* 2007;62(4):278-83.  
<http://www.ncbi.nlm.nih.gov/pubmed/17484283>
5. Duchene D, Wouessidjewe D. Pharmaceutical uses of cyclodextrins and derivatives. *Drug Dev. Ind. Pharm.* 1990;16(17):2487-99.  
<http://informahealthcare.com/doi/abs/10.3109/03639049009058543>
6. Indian Pharmacopoeia, Ministry of Health and Family Welfare, Controller of Publication, New Delhi: 1996.
7. Mura P, Maestrelli F, Cirri M. Ternary systems of naproxen with hydroxypropyl-beta-cyclodextrin and aminoacids. *Int. J. Pharm.* 2003;260(2):293-302.  
<http://www.ncbi.nlm.nih.gov/pubmed/12842348>
8. Naidu NB, Chowdary KP, Murthy KV, Satyanarayana V, Hayman AR, Becket G. Physicochemical characterization and dissolution properties of meloxicam-cyclodextrin binary systems. *J. Pharm. Biomed. Anal.* 2004;35(1):75-86.  
<http://www.ncbi.nlm.nih.gov/pubmed/15030882>
9. Ribeiro LS, Ferreira DC, Veiga FJ. Physicochemical investigation of the effects of water-soluble polymers on vinpocetine complexation with beta-cyclodextrin and its sulfobutyl ether derivative in solution and solid state. *Eur. J. Pharm. Sci.* 2003;20(3):253-66.  
<http://www.ncbi.nlm.nih.gov/pubmed/14592691>

10. Sachan NK, Pushkar S. Solid dispersions: An industrially feasible alternative approach to formulate brick dust molecules. *Bull. Pharm. Res.* 2011;1(1):75-87.  
<http://www.appconnect.in/wp-content/uploads/2012/01/ReprintBPR013.pdf>
11. Sami F, Philip B, Pathak K. Effect of auxiliary substances on complexation efficiency and intrinsic dissolution rate of gemfibrozil-beta-CD complexes. *AAPS PharmSciTech.* 2010;11(1):27-35.  
<http://www.ncbi.nlm.nih.gov/pmc/articles/PMC2850474/>
12. complexes of glyburide with hydroxypropyl- $\beta$ -cyclodextrin and other solubilizing agents and their effect on release behavior of glyburide in aqueous and buffered media at different agitation speeds. *Drug Dev. Ind. Pharm.* 2012;38(11):1328-36.  
<http://www.ncbi.nlm.nih.gov/pubmed/22283512>
13. Taupitz T, Dressman JB, Buchanan CM, Klein S. Cyclodextrin-water soluble polymer ternary complexes enhance the solubility and dissolution behaviour of poorly soluble drugs. Case example: itraconazole. *Eur. J. Pharm. Biopharm.* 2013;83(3):378-87.  
<http://www.ncbi.nlm.nih.gov/pubmed/23201048>
14. Wang D, Li H, Gu J, Guo T, Yang S, Guo Z, Zhang X, Zhu W, Zhang J. Ternary system of dihydroartemisinin with hydroxypropyl- $\beta$ -cyclodextrin and lecithin: simultaneous enhancement of drug solubility and stability in aqueous solutions. *J. Pharm. Biomed. Anal.* 2013;83:141-8.  
<http://www.sciencedirect.com/science/article/pii/S0731708513001945>

