

Agarwal S, Mishra VK, Mishra M, Singh S, Kumar V, Kashaw V, Kashaw SK. QSAR studies involving 2D, 3D QSAR and pharmacophore mapping studies on arylsulfonyl imidazolidinone derivatives as anticancer agents. *Bull. Pharm. Res.* 2015;5(2):70-80.

References (18):

1. Abdalla A, Klein S, Mader K. A new self emulsifying drug delivery system for poorly soluble drugs: characterization, dissolution, in vitro digestion and incorporation into solid pellets. *Eur. J. Pharm. Sci.* 2008;35(5):457-464. [DOI: 10.1016/j.ejps.2008.09.006]
<http://www.ncbi.nlm.nih.gov/pubmed/18940249>
2. Amidon GL, Lennernas H, Shah VP, Crison JR. A theoretical basis for a biopharmaceutic drug classification: the correlation of in vitro drug product dissolution and in vivo bioavailability. *Pharm. Res.* 1995;12(3):413-20.
<http://www.ncbi.nlm.nih.gov/pubmed/7617530>
3. Craig DQM, Barker SA, Banning D, Booth SW. An investigation into the mechanisms of self-emulsification using particle size analysis and low frequency dielectric spectroscopy. *Int. J. Pharm.* 1995;114(1):103-10.
<http://www.sciencedirect.com/science/article/pii/0378517394002220>
4. Dahiya S, Tayde P. Binary and ternary solid systems of carvedilol with 2-hydroxypropyl- β -cyclodextrin and PVP K30. *Bull. Pharm. Res.* 2013;3(3):128-34.
<http://journal.appconnect.in/wp-content/uploads/2013/10/ReprintBPR082.pdf>
5. Gershanik T, Benita S. Self-dispersing lipid formulations for improving oral absorption of lipophilic drugs. *Eur. J. Pharm. Biopharm.* 2000;50(1):179-88.
<http://www.ncbi.nlm.nih.gov/pubmed/10840200>
6. Gursoy RN, Benita S. Self-emulsifying drug delivery systems (SEDDS) for improved oral delivery of lipophilic drugs. *Biomed. Pharmacother.* 2004;58(3):173-82.
<http://www.sciencedirect.com/science/article/pii/S0753332204000319>
7. Horter D, Dressman JB. Influence of physicochemical properties on dissolution of drugs in the gastrointestinal tract. *Adv. Drug Deliv. Rev.* 2001;46(1-3):75-87.
<http://www.sciencedirect.com/science/article/pii/S0169409X00001307>
8. Nazzal S, Guven N, Reddy IK, Khan MA. Preparation and characterization of coenzyme Q10-Eudragit solid dispersion. *Drug Dev. Ind. Pharm.* 2002;28(1):49-57.
<http://www.ncbi.nlm.nih.gov/pubmed/11858524>

9. Pabreja K, Dua K. Comparative evaluation of *in situ* intestinal absorption of aceclofenac from solid dispersions, β -cyclodextrin complexes and co-precipitates in rats. *Bull. Pharm. Res.* 2011;1(1):26-30.
<http://www.appconnect.in/wp-content/uploads/2012/01/ReprintBPR005.pdf>
10. Patel AR, Vavia PR. Preparation and in vivo evaluation of SMEDDS (self-microemulsifying drug delivery system) containing fenofibrate. *AAPS J.* 2007;9(3):E344-52.
<http://www.ncbi.nlm.nih.gov/pubmed/18170981>
11. Prusty A. Formulation and in-vitro evaluation of amlodipine besylate-HP- β -CD inclusion complex incorporated mouth dissolving tablets. *Bull. Pharm. Res.* 2014;4(3):124-8.
<http://journal.appconnect.in/wp-content/uploads/2015/01/Reprintbpr102.pdf>
12. Sahoo DK, Sahu PK, Patro CS. New validated isocratic RP-HPLC method for assay of fenofibrate. *Int. J. Pharm. Pharm. Sci.* 2014;6(2):169-172.
<http://www.ijppsjournal.com/Vol6Suppl2/7918.pdf>
13. Shah NH, Carvajal MT, Patel CI, Infeld MH, Malick AW. Self-emulsifying drug delivery systems (SEDDS) with polyglycolized glycerides for improving in vitro dissolution and oral absorption of lipophilic drugs. *Int J Pharm.* 1994;106(1):15-23.
<http://www.sciencedirect.com/science/article/pii/0378517394902712>
14. Shah SP, Shah MD, Agrawal YK. Self-micro emulsifying drug delivery system: a novel approach for enhancement of oral bioavailability of poorly soluble drugs. *Am. J. PharmTech Res.* 2012;2(1):193-215.
file:///C:/Users/Home/Downloads/AJPTR%20article%20sanket_8630.pdf
15. Shen H, Zhong M. Preparation and evaluation of self-microemulsifying drug delivery systems (SMEDDS) containing atorvastatin. *J. Pharm. Pharmacol.* 2006;58(9):1183-91.
<http://www.ncbi.nlm.nih.gov/pubmed/16945176>
16. Stegemann S, Leveiller F, Franchi D, de Jong H, Linden H. When poor solubility becomes an issue: from early stage to proof of concept. *Eur. J. Pharm. Sci.* 2007;31(5):249-61.
<http://www.ncbi.nlm.nih.gov/pubmed/17616376>
17. Sugimoto M, Okagaki T, Narisawa S, Koida Y, Nakajima K. Improvement of dissolution characteristics and bioavailability of poorly water-soluble drugs by novel cogrinding method using water-soluble polymer. *Int. J. Pharm.* 1998;160(1):11-9.
<http://www.sciencedirect.com/science/article/pii/S0378517397002937>
18. Tang B, Cheng G, Gu JC, Xu CH. Development of solid self-emulsifying drug delivery systems: preparation techniques and dosage forms. *Drug. Discov. Today* 2008;13(13-14):606-12.
<http://www.ncbi.nlm.nih.gov/pubmed/18598917>