

REFERENCES

- Agarwal A, Bora D, Agarwal C, Kumar R, Choudhary V. CNS stimulant and antidepressant activity of seeds of *Abelmoschus esculentus* in rats. *Bull. Pharm. Res.* 2015; 5(2):47-50.
- Costa SCC, Detoni CB, Branco CRC, Botura MB, Branco A. In vitro photoprotective effects of *Marcetia taxifolia* ethanolic extract and its potential for sunscreen formulations. *Brazil. J. Pharmacogn.* 2015;25(4):413-8. [DOI: 10.1016/j.bjp.2015.07.013]
- Degirmenci NS, Akal ZU, Alpsoy L. Cytotoxic and apoptotic effects of caffeic acid and caffeoyl malic acid on brain glioblastoma (U87-MG). *Bull. Pharm. Res.* 2016;6(1):7-13. [DOI: 10.21276/bpr.2016.6.1.2]
- Dahiya R. Synthesis of a phenylalanine-rich peptide as potential anthelmintic and cytotoxic agent. *Acta Pol. Pharm.* 2007;64(6):509-16.
- Dahiya R. Total synthesis and biological potential of psammosilenin A. *Arch. Pharm. (Weinheim)* 2008;341(8): 502-9. [DOI: 10.1002/ardp.200800006]
- Dahiya R, Kumar A. Synthetic and biological studies on a cyclopolypeptide of plant origin. *J. Zhejiang Univ. Sci. B* 2008;9(5):391-400. [DOI: 10.1631/jzus.B0720001]
- Dahiya R, Singh S. Synthesis, characterization, and biological activity studies on fanlizhicyclopeptide A. *Iran. J. Pharm. Res.* 2017;16(3):1176-84.
- Dahiya R, Singh S, Kaur K, Kaur R. Total synthesis of a natural cyclooligopeptide from fruits of sugar-apples. *Bull. Pharm. Res.* 2017;7(3):151. [DOI: 10.21276/bpr.2017.7.3.4]
- Ellman GL. Tissue sulfhydryl group. *Arch. Biochem. Biophys.* 1959;82(1):70-7. [DOI: 10.1016/0003-9861(59)90090-6]
- Emran TB, Rahman MA, Zahid Hosen SM, Saha D, Chowdhury S, Saha D, Dey TK. Antioxidant property of ethanolic extract of *Leucas aspera* Linn. *Bull. Pharm. Res.* 2012;2(1):46-9.
- Fulton JE Jr. Utilizing the ultraviolet (UV detect) camera to enhance the appearance of photodamage and other skin conditions. *Dermatol. Surg.* 1997;23(3):163-9. [DOI: 10.1111/j.1524-4725.1997.tb00013.x]
- Gajendiran A, Thangaraman V, Thangamani S, Ravi D, Abraham J. Antimicrobial, antioxidant and anticancer screening of *Ocimum basilicum* seeds. *Bull. Pharm. Res.* 2016;6(3):114-9. [DOI: 10.21276/bpr.2016.6.3.5]
- Grant K. Top-selling herbal supplements. *J. Manag. Care Spec. Pharm.* 1999;5(4):357-70. [DOI: 10.18553/jmcp.1999.5.4.357]
- Jung JW, Cha SH, Lee SC, Chun IK, Kim YP. Age related changes of water content in the rate skin. *J. Dermatol. Sci.* 1997;14(1):12-9. [DOI: 10.1016/S0923-1811(96)00542-7]
- Kim BJ, Kim JH, Kim HP, Heo MY. Biological screening of 100 plant extracts for cosmetic use (II): anti-oxidative activity and free radical scavenging activity. *Int. J. Cosmet. Sci.* 1997;19(6):299-307. [DOI: 10.1046/j.1467-2494.1997.171726.x]
- Kim SJ, Lim MH, Chun IK, Won YH. Effect of flavonoids of *Ginkgo biloba* on proliferation of human skin fibroblast. *Skin Pharmacol.* 1997;10(4):200-5. [DOI: 10.1159/00021505]
- Kligman LH. Photoaging. Manifestations, prevention and treatment. *Clin. Geriatr. Med.* 1989;5(1):235-51.
- Korac RR, Khambholja KM. Potential of herbs in skin protection from ultraviolet radiation. *Pharmacogn. Rev.* 2011;5(10):164-73. [DOI: 10.4103/0973-7847.91114]
- Milbury PE, Chen C-Y, Dolnikowski GG, Blumberg JB. Determination of flavonoids and phenolics and their distribution in almonds. *J. Agri. Food. Chem.* 2006;54(14): 5027-33. [DOI: 10.1021/jf0603937]
- Monagas M, Garrido I, Lebron-Aguilar R, Bartolome B, Gomez-Cordoves C. Almonds (*Prunus dulcis* (Mill.) D.A. Webb) skin as a potential source of bioactive polyphenols. *J. Agric. Food Chem.* 2007;55(21):8498-507. [DOI: 10.1021/jf071780z]
- Nicol NJ, Fenske NA. Photo damage: causes, clinical manifestations, and prevention. *Dermatol. Nurs.* 1993; 5(4):263-75.
- Ohkawa H, Ohishi N, Yagi K. Assay for lipid peroxides in animal tissues by thiobarbituric acid reaction. *Anal. Biochem.* 1979;95(2):351-8. [DOI: 10.1016/0003-2697(79)90738-3]
- Oloke J, Odelade K, Oladeji O. Characterization and antimicrobial analysis of flavonoids in *Vernonia amygdalina*: a common chewing stick in south-western Nigeria. *Bull. Pharm. Res.* 2017;7(3):149. [DOI: 10.21276/bpr.2017.7.3.2]
- Ryu A, Naru E, Arakane K, Masunaga T, Shinmoto K, Nagano T, Hirobe M, Mashiko S. Cross linking of collagen by singlet oxygen generated with UV-A. *Chem. Pharm. Bull. (Tokyo)* 1997;45(8):1243-7.
- Senthil Kumar R, Vinoth Kumar S, Sudhakar P. Anticancer activity of methanolic leaf extract of *Morinda tinctoria* roxb. against ehrlich ascites carcinoma in mice. *Bull. Pharm. Res.* 2017;7(2):146. [DOI: 10.21276/bpr.2017.7.2.4]
- Shrestha DK, Sapkota H, Baidya P, Basnet S. Antioxidant and antibacterial activities of *Allium sativum* and *Allium cepa*. *Bull. Pharm. Res.* 2016;6(2):50-5. [DOI: 10.21276/bpr.2016.6.2.3]
- Stahl W, Sies H. Carotenoids and flavonoids contribute to nutritional protection against skin damage from sunlight. *Mol. Biotechnol.* 2007;37(1):26-30.
- Steenvoorden DPT, van Henegouwen GMJB. The use of endogenous antioxidants to improve photoprotection. *J. Photochem. Photobiol. B: Biol.* 1997;41(1-2):1-10. [DOI: 10.1016/S1011-1344(97)00081-X]
- Taylor CR, Stern RS, Leyden JJ, Gilchrist BA. Photoaging/ photodamage and photoprotection. *J. Am. Acad. Dermatol.* 1990;22(1):1-15. [DOI: 10.1016/0190-9622(90)70001-X]
- Viana CB, Carbonezi LH, Martins RCC. Isolation of pentacyclic triterpenes from *Simira sampaioana* (standl.) steyererm (Rubiaceae) as possible anticancer agents. *Bull. Pharm. Res.* 2017;7:142. [DOI: 10.21276/bpr.2017.7.1.5]
