

Patel TB, Patel LD, Patel TR, Suhagia BN. Artificial neural network as tool for quality by design in formulation development of solid dispersion of fenofibrate. *Bull. Pharm. Res.* 2015;5(1):20-7.

### References (27):

1. Agatonovic-Kustrin S, Beresford R (2000) Basic concepts of artificial neural network (ANN) modeling and its application in pharmaceutical research. *J. Pharm. Biomed. Anal.* 2000;22(5):717-27.  
<http://www.ncbi.nlm.nih.gov/pubmed/10815714>
2. Ahmadi Lakalayeh G, Faridi-Majidi R, Saber R, Partoazar A, Mehr SE, Amami A. Investigating the parameters affecting the stability of superparamagnetic iron oxide-loaded nanoemulsion using artificial neural networks. *AAPS PharmSciTech.* 2012;13(4):1386-95.  
<http://www.ncbi.nlm.nih.gov/pubmed/23054990>
3. Aksu B, Paradkar A, de Matas M, Ozer O, Guneri T, York P. Quality by design approach: application of artificial intelligence techniques of tablets manufactured by direct compression. *AAPS PharmSciTech.* 2012;13(4):1138-46.  
<http://www.ncbi.nlm.nih.gov/pubmed/22956056>
4. Anand P, Siva Prasad BVN, Venkateswarlu CH. Modeling and optimization of a pharmaceutical formulation system using radial basis function network. *Int. J. Neural. Syst.* 2009;19(2):127-36.  
<http://www.ncbi.nlm.nih.gov/pubmed/19496208>
5. Arulsudar N, Subramanian N, Muthy RSR. Comparison of artificial neural network and multiple linear regression in the optimization of formulation parameters of leuprolide acetate loaded liposomes. *J. Pharm. Pharmaceut. Sci.* 2005;8(2):243-58.
6. Barmpalexis P, Kanaze FI, Kachrimanis K, Georgarakis E. Artificial neural networks in the optimization of a nimodipine controlled release tablet formulation. *Eur. J. Pharm. Biopharm.* 2010;74:316-23.  
<http://www.ncbi.nlm.nih.gov/pubmed/19815063>
7. Bourquin J, Schmidli H, van Hoogevest P, Leuenberger H. Advantages of artificial neural networks (ANNs) as alternative modelling technique for data sets showing non-linear relationships using data from a galenical study on a solid dosage form. *Eur. J. Pharm. Sci.* 1998;7(1):5-16.  
<http://www.ncbi.nlm.nih.gov/pubmed/9845773>

8. Gasperlin M, Podlogar F, Sibanc R. Evolutionary artificial neural networks as tools for predicting the internal structure of microemulsions. *J. Pharm. Pharmaceut. Sci.* 2008;11(1): 67-76.  
<http://ejournals.library.ualberta.ca/index.php/JPPS/article/view/1570/1096>
9. Hussain AS, Yu XQ, Johnson RD. Application of neural computing in pharmaceutical product development. *Pharm. Res.* 1991;8(10):1248-52.  
<http://www.ncbi.nlm.nih.gov/pubmed/1796042>
10. Husseini GA, Abdel-Jabbar NM, Mjalli FS, Pitt WG. Modeling and sensitivity analysis of acoustic release of Doxorubicin from unstabilized pluronic P105 using an artificial neural network model. *Technol. Cancer Res. Treat.* 2007;6(1):49-56.  
<http://www.ncbi.nlm.nih.gov/pubmed/17241100>
11. Ibrić S, Jovanović M, Djurić Z, Parojčić J, Petrović SD, Solomun L, Stupar B. Artificial neural networks in the modeling and optimization of aspirin extended release tablets with Eudragit L 100 as matrix substance. *AAPS PharmSciTech.* 2003;4(1):62-70.  
<http://www.ncbi.nlm.nih.gov/pmc/articles/PMC2750305/>
12. Ioele G, De Luca M, Dinç E, Oliverio F, Ragno G. (2011) Artificial neural network combined with principal component analysis for resolution of complex pharmaceutical formulations. *Chem. Pharm. Bull. (Tokyo)* 2011;59(1):35-40.  
<http://www.ncbi.nlm.nih.gov/pubmed/21212544>
13. Ivić B, Ibrić S, Cvetković N, Petrović A, Trajković S, Djurić Z. Application of design of experiments and multilayer perceptrons neural network in the optimization of diclofenac sodium extended release tablets with Carbopol® 71G. *Chem. Pharm. Bull. (Tokyo)* 2010;58(7):947-9.  
<http://www.ncbi.nlm.nih.gov/pubmed/20606343>
14. Jouyban A, Majidi M-R, Jalilzadeh H, Asadpour-Zeynali K. Modeling drug solubility in water-cosolvent mixtures using an artificial neural network. *Il Farmaco* 2004;59(6):505-12.  
<http://www.sciencedirect.com/science/article/pii/S0014827X04000576>
15. Kikuchi S, Takayama K. Reliability assessment for the optimal formulations of pharmaceutical products predicted by a nonlinear response surface method. *Int. J. Pharm.* 2009;374(1-2):5-11.  
<http://www.ncbi.nlm.nih.gov/pubmed/19446752>
16. Kikuchi S, Takayama K. Multivariate statistical approach to optimizing sustained-release tablet formulations containing diltiazem hydrochloride as a model highly water-soluble drug. *Int. J. Pharm.* 2010;386(1-2):149-55.  
<http://www.ncbi.nlm.nih.gov/pubmed/19922778>

17. Leane MM, Cumming I, Corrigan OI. The use of artificial neural networks for the selection of the most appropriate formulation and processing variables in order to predict the in vitro dissolution of sustained release minitables. *AAPS PharmSciTech*. 2003;4(2):129-40.  
<http://www.ncbi.nlm.nih.gov/pmc/articles/PMC2750588/>
18. Mandal U, Gowda V, Ghosh A, Bose A, Bhaumik U, Chatterjee B, Pal TK. Optimization of metformin HCl 500 mg sustained release matrix tablets using artificial neural network (ANN) based on Multilayer Perceptrons (MLP) model. *Chem. Pharm. Bull. (Tokyo)* 2008; 56(2):150-5.  
<http://www.ncbi.nlm.nih.gov/pubmed/18239298>
19. Mendyk A, Jachowicz R, Doroczyński P. Artificial neural networks in the modeling of drugs release profiles from hydrodynamically balanced systems. *Acta Pol. Pharm.* 2006;63(1):75-80.  
[http://www.ptfarm.pl/pub/File/Acta\\_Poloniae/2006/1/75.pdf](http://www.ptfarm.pl/pub/File/Acta_Poloniae/2006/1/75.pdf)
20. Mendyk A, Tuszyński PK, Polak S, Jachowicz R (2013) Generalized in vitro-in vivo relationship (IVIVR) model based on artificial neural networks. *Drug Des. Devel. Ther.* 2013;7:223-32.  
<http://www.ncbi.nlm.nih.gov/pubmed/23569360>
21. Nelofer R, Ramanan RN, Rahman RN, Basri M, Ariff AB. Comparison of the estimation capabilities of response surface methodology and artificial neural network for the optimization of recombinant lipase production by E. coli BL21. *J. Ind. Microbiol. Biotechnol.* 2012;39(2):243-54.  
<http://www.ncbi.nlm.nih.gov/pubmed/21833714>
22. Nettles DL, Haider MA, Chilkoti A, Setton LA. Neural network analysis identifies scaffold properties necessary for in vitro chondrogenesis in elastin-like polypeptide biopolymer scaffolds. *Tissue Eng. Part A* 2010;16(1):11-20.  
<http://www.ncbi.nlm.nih.gov/pubmed/19754250>
23. Sadatsafavi M, Moayyeri A, Soltani A, Larijani B, Nouraie M, Akhondzadeh S. Artificial neural networks in prediction of bone density among post-menopausal women. *J. Endocrinol. Invest.* 2005;28(7):425-31.  
<http://www.ncbi.nlm.nih.gov/pubmed/16075926>
24. Sibanda W, Pillay V, Danckwerts MP, Viljoen AM, van Vuuren S, Khan RA. Experimental design for the formulation and optimization of novel cross-linked oilispheres developed for in vitro site-specific release of Mentha piperita oil. *AAPS PharmSciTech*. 2004;5(1):128-41.  
<http://www.ncbi.nlm.nih.gov/pmc/articles/PMC2784852/>
25. Takayama K, Fujikawa M, Nagai T. Artificial neural network as a novel method to optimize pharmaceutical formulations. *Pharm. Res.* 1999;16(1):1-6.  
<http://link.springer.com/article/10.1023%2FA%3A1011986823850>

26. Takayama K, Morva A, Fujikawa M, Hattori Y, Obata Y, Nagai T. Formula optimization of theophylline controlled-release tablet based on artificial neural networks. *J. Control. Release* 2000;68(2):175-86.  
<http://www.ncbi.nlm.nih.gov/pubmed/10925126>
27. Takayama K, Obata Y, Morishita M, Nagai T. Multivariate spline interpolation as a novel method to optimize pharmaceutical formulations. *Pharmazie* 2004;59(5):392-5.  
<http://www.ncbi.nlm.nih.gov/pubmed/15212308>

