



RESEARCH PAPER

EFFECTS OF COMBINED EXPOSURE TO DICHLORVOS AND MONOCROTOPHOS ON HEPATOTOXICITY IN RATS

Vinesh Kumar^{1,2*}, Pawan Kumar Basniwal² and R. Vijayaraghavan¹

¹Division of Pharmacology and Toxicology, Defense Research and Development Establishment (DRDE), Gwalior- 474002, Madhya Pradesh, India

²Department of Pharmaceutical Chemistry, Lal Bahadur Shastri College of Pharmacy, Jaipur-302004, Rajasthan, India

*E-mail: vineshkc@gmail.com

Tel.: +91 9828262634.

Received: Mar 07, 2018 / Revised: Apr 11, 2018 / Accepted: Apr 12, 2018

Blind use of organophosphates as pesticides for enhancing productivity in agriculture leads to hormonal imbalance, infertility, thyroid dysfunctioning and the degree of effects produced depends on the extent of exposure to pesticides. Thus, the present study was planned to determine the effect of individual and combined exposure to dichlorvos (DDVP) and monocrotophos (MCP) on some selected biochemical variables suggestive of hepatic damage in rats. The monocrotophos clearly possessed the propensity to exacerbate hepatotoxicity when administered even at a relatively low dose, while the exact mechanisms by which monocrotophos potentiate the toxicity needs further investigation. The possibility of oxidative stress in the augmentation of hepatotoxicity was investigated that MCP is more potent in inducing oxidative stress. Therefore, the present study provides some interesting new observations for possible co-exposure to organophosphates. The co-exposure to DDVP and MCP may produce synergistic effects at some extent.

Key words: Dichlorvos, Monocrotophos, Hepatotoxicity, Rats, Pesticide, Organophosphate.

INTRODUCTION

Pesticides have been used in agriculture for centuries to increase food production by removing unwanted insects and controlling disease vectors (Environmental Protection Agency US, 2000). Around four million tons of pesticides are applied to crops annually for pest control over the world, but less than one percent of the total applied pesticides reach the target pests (Pimentel, 1983). This massive use causes run off of pesticides from agricultural fields leading to contamination and bioaccumulation in tissues of many species resulting in development of toxicity (Elia *et al* 2006).

These pesticides are unselective and toxic to wide variety of species including humans (Dawson *et al* 2010). Poisoning occurs as a result of agricultural use, accidental exposure, suicide

and, rarely homicidal use (Aygun, 2004).

Consumption of contaminated food items with pesticides above maximum residue level results in development of short term and long term adverse effects. It is evident from the incident in which many people died after consumption of pesticide contaminated flour (Gupta, 2004).

Action of pesticides on different body organs affects normal body functioning and causes problems like hepatitis, degeneration of liver, dyspnea and burning sensation in urine (Azmi *et al* 2006; Hettwer, 1975). Exposure to pesticides can affect hormonal balance, reproduction, thyroid functioning (Tyler *et al* 1998; Ewing, 1999). Degree of effects produced depends on the extent of exposure to pesticides (Chitra *et al* 2006). Organophosphorus (OP) toxicity is one of the major health issues globally and it results

around 200,000 deaths per year all over the world due to self ingestion of these compounds (Eddleston, 2008). Oxidative damage is thought to be an important mechanism of damage in organophosphate pesticides toxicity (Banerjee *et al* 2001; Delescluse *et al* 2001; Halliwell *et al* 2004; Dwivedi *et al* 2011). Organophosphate pesticides have also been reported to reduce antioxidant enzyme activity, enhance the production of lipid peroxides and reduce the level of cellular antioxidants (Julka *et al* 1992). Pesticides, especially organophosphate pesticides, induce oxidative stress both *in vivo* and *in vitro* (Bagchi *et al* 1995; Yang *et al* 1998). Dichlorvos, also known as DDVP or DDVF (2, 2-dichlorovinyl dimethyl phosphate) is a highly volatile, directly acting acetylcholine esterase inhibitor (Environmental Protection Agency US, 2000). The irreversible binding to and subsequent inactivation of acetylcholine esterase, the enzyme that normally catalyzes the hydrolysis of acetylcholine (ACh) at neuromuscular junctions and other cholinergic synapses, is generally believed to be the major mechanism of their toxicity. The subsequent accumulation of ACh in the cholinergic clefts causes overstimulation of the peripheral as well as the central cholinergic nervous system resulting in clinical manifestation in the form of acute cholinergic crisis (Taylor, 1996).

Another widely used pesticide, Monocrotophos (dimethyl (E)-1-methyl-2-(methylcarbamoyl) vinyl phosphate) is a broad spectrum systemic insecticide and acaricide used on variety of crops such as cotton, rice, and sugarcane. Exposure of monocrotophos is known to produce a variety of biochemical changes in mammals, aquatic and other experimental animals (Nemcsok *et al* 1987; Sultatos *et al* 1994; Waite *et al* 1992). It is a systemic pesticide harmful to human beings, affecting eyes and the CNS (Horrigan *et al* 2002). Monocrotophos along with endosulfan interferes with NADPH dependent monooxygenase mechanism and is effective inducers of NADPH cytochrome C reductase (Ramaneswari *et al* 2008).

Both DDVP and MCP possess different characteristics and these compounds produce varying degree of toxic effects. Liver is known to be the major target organ for pesticides, and it is an important organ for metabolic waste excretion and pesticide elimination (Shugart *et al* 1992). Analysis of biochemical parameters could help to identify target organs of toxicity. It may also provide an early warning signal in

stressed organism (Folmar, 1993). OPs have been reported to cause enzyme induction particularly of ALP, ALT, AST, GGT, and LDH (Kalender *et al* 2005). Changes in the levels of these enzymes may differ depending on the exposure time and dose of OP. It has been previously suggested that organophosphates may phosphorylate and inhibit the hydroxymethylglutaryl CoA reductase, the key enzyme in cholesterol production (Ryhanen *et al* 1984). Increase in the activities of the liver specific enzymes, transaminases, alanine aminotransferase (ALT) and aspartate aminotransferase (AST) in blood have been attributed to tissue damage, particularly the liver and are considered as diagnostic and sensitive markers of hepatic damage (Reichling *et al* 1988). Hence, measurement of transaminase activities in plasma has been used as an indicator of hepatotoxicity of pesticides (Agrahari *et al* 2007). AST is an enzyme that is normally present in liver and heart cells. The blood AST levels are thus elevated with liver damage or with an insult to the heart. While, as a liver specific enzyme, ALT is significantly elevated in conditions of hepatic damage although increases can occur in connection with damage of heart as well. Hence, AST/ALT ratio is often employed for the differential diagnosis of liver damage, with the higher ratio indicating more severe damage.

Various studies have been carried out on individual effects of these two organophosphates but there are only few studies on combined exposure of DDVP and MCP (Dwivedi *et al* 2010; Dwivedi *et al* 2011). Considering the fact that different pesticides are simultaneously being used, there is every possibility of population getting exposed to multiple pesticides. Thus, the present study was planned to determine the effect of individual and combined exposure to DDVP and MCP on some selected biochemical variables suggestive of hepatic damage in rats.

EXPERIMENTAL

Chemicals

The two organophosphorous compounds, Dichlorvos (Nuvan, 76%) and MCP (Kadett, 36%) were obtained from Syngenta chemicals and P.I. Industries Ltd, respectively. All other chemicals and reagents were of analytical grade. All other analytical laboratory chemicals and reagents were purchased from Merck (Germany), Sigma (USA) or BDH chemicals (Mumbai, India). Ultra pure water prepared by Millipore (New Delhi, India) was used throughout the experiment to

avoid metal contamination and for the preparation of reagents and buffers used for various biochemical assays in our study.

Animals

Wistar rats (110–120 g) were obtained from Defence Research and Development Establishment (DRDE) animal facility and prior to use, were acclimatized for 7 days 12 h light/dark cycle. The animal ethical committee of DRDE, Gwalior approved the protocols for the experiments. The animals were housed in stainless steel cages in an air-conditioned room with temperature maintained at 25±2°C. Rats were allowed standard pellet diet (Ashirwad Feeds, Chandigarh, India) throughout the experiment and water *ad libitum*. Different doses (high dose and low dose) of DDVP and MCP were selected to study the effects on blood

biochemical variables in experimental animals (**Table 1**).

Biochemical assays

Blood and tissue lipid peroxidation was measured (Ohkawa *et al* 1979) and modified (Saxena *et al* 2004). Reactive oxygen species (ROS) assay was performed (Socci *et al* 1999). The activity of SGOT and SGPT were assayed according to the literature method (Reitman *et al* 1957). Reduced glutathione (GSH) and oxidized glutathione (GSSG) levels were measured fluorometrically (Hissin *et al* 1974). Catalase activity in tissue was assayed following the procedure reported in literature (Sinha, 1972). SOD activity was assayed and total tissue protein was measured by the reported methods (Nishikimi *et al* 1972; Kakkar *et al* 1984; Lowry *et al* 1951).

Table 1. Dosing of the groups of animals

Group	Number of animals	Drug, dose and route	
		High dose group	Low dose group
Group 1	Six	Drinking water (blank)	Drinking water (blank)
Group 2	Six	Dichlorvos, 2.5 mg/kg, subcutaneously	Dichlorvos, 2.0 mg/kg, orally
Group 3	Six	Monocrotophos, 2.0 mg/kg, orally	Monocrotophos, 1.8 mg/kg, orally
Group 4	Six	Dichlorvos, 2.5 mg/kg, subcutaneously + Monocrotophos, 2.0 mg/kg, orally	Dichlorvos, 1 mg/kg orally + Monocrotophos, 1 mg/kg orally

Statistical analysis

The results are expressed as the mean±SEM of number of observations. Comparisons of means were carried out using one way ANOVA followed by student 't' test to compare means between the different treatment groups. Differences were considered significant at P<0.05 unless otherwise stated in the text.

RESULTS

Effects at high dose

Effects on biochemical variables indicative of liver damage

Effects of individual and combined exposure to DDVP and MCP on biochemical variables indicative of hepatic damage are presented in the **Table 2**. Exposure to MCP alone led to a significant increase in serum AST and ALP activities suggesting hepatic injury while ALT activity decreased significantly. Exposure to DDVP produced no change in these variables.

Combined exposure to DDVP and MCP had no additional synergistic additive effects however; the increased AST activity during combined exposure was predominantly due to MCP.

Effects at low dose

Hepatic ROS and TBARS levels

Effect of DDVP and MCP on some hepatic biochemical variables is shown in **Figure 1**. Co-exposure to DDVP and MCP produced a non-significant elevation in ROS level. Interestingly, exposure to DDVP and MCP led to a significant elevation in TBARS level both in alone and combination groups (**Figure 3**).

Hepatic GSH and GSSG levels

Effect of DDVP and MCP on hepatic GSH and GSSG levels is shown in **Figure 2**. GSH and GSSG levels did not found to alter following exposure to MCP and DDVP whether administered alone or in combination.

Table 2. Effects of individual and combined exposure of DDVP and MCP on hepatic ALT and AST levels in rats at high dose

Variables	Normal	DDVP	MCP	DDVP + MCP
S-AST (U/L)	35.85±1.76*	33.60±1.06*	138.45±9.93†	134.58±9.09†
S-ALT (U/L)	73.59±4.28*	73.41±4.40*	67.93±3.60*	78.24±7.30*
ALP (mg of P/hr/mg protein)	0.13±0.05*	0.16±0.02*	0.18±0.04*	0.19±0.03*


S-AST- Serum aspartate amino transferase; S-ALT- Serum alanine amino transferase; ALP- Alkaline phosphatase, values are mean±SE; n=5; Data was analyzed using analysis of variance (ANOVA) followed by Dunnett's test. *†Differences between values with matching symbol notations within each row are not statistically significant at a 5% level of probability

Hepatic SOD and catalase activities

The activity of SOD decreased significantly in MCP exposed group as well as in the combination group. However, the activity of catalase was found to be decreased in DDVP and combination group (**Figure 3**).


SGOT and SGPT activity

The activities of SGOT and SGPT indicative of liver damage in rats are depicted in **Figure 4**. Serum SGOT and SGPT activities increased on exposure to DDVP and MCP individually and on co-exposure suggesting liver injury.



ROS, reactive oxygen species as FIU; TBARS, thiobarbituric acid reactive species as mg/gm tissue; values are mean±SE; n = 5 *†Differences between values with matching symbol notations within each column are not statistically significant at a 5% level of probability

Fig. 1. Effect of individual and combined exposure to DDVP and MCP on ROS and TBARS in rat liver at low dose



GSH, reduced glutathione as mg/gm tissue; GSSG, oxidized glutathione as mg/gm tissue; values are mean±SE; n = 5 *†Differences between values with matching symbol notations within each column are not statistically significant at a 5% level of probability

Fig. 2. Effect of individual and combined exposure to DDVP and MCP on GSH and GSSG in rat liver low dose




Fig. 3. Effect of individual and combined exposure to DDVP and MCP on SOD and Catalase in rat liver at low dose




Fig. 4. Effect of individual and combined exposure to DDVP and MCP on SGOT and SGPT in rats at low dose

DISCUSSION

Organophosphorus pesticides being lipophilic interact with cells through lipid-rich bio-membranes and damage membrane as a result of oxidative polyunsaturated fatty acids of the bilayer known as lipid per oxidation (Mittal *et al* 2006; Naqvi *et al* 1992; Yamano *et al* 1992). Transaminases (SGOT and SGPT) are critical enzymes in the biological processes. Increased activity of transaminases is mainly due to the leakage of these enzymes from liver cytosol into the blood stream. Thus, their increased activity in serum is an indicative of liver damage. Higher dose of both the toxicants produced liver injury and the impairment in liver function was supported by the liver function test observations. Exposure to DDVP or MCP led to hepatic damage indicated by increased SGOT and

SGPT activities in serum. There was however more pronounced increase in SGOT activity in animals co-exposed to DDVP and MCP suggesting their cumulative effect (Dwivedi *et al* 2010). Since, higher dose of DDVP and MCP were capable enough of producing pronounced toxicity, a lower dose toxicity study was carried out to investigate changes in oxidative stress which may be a potential cause of toxicity. Earlier reports have investigated toxic effects of individual exposure of DDVP and MCP in experimental animals. However, there are only few studies in the literature which reported combined toxicity of these toxicants (Dwivedi *et al* 2011). There is paucity of evidence as the interactions between these two OP's have not been investigated in detail. The present study thus, observed the effects of combined exposure

of DDVP and MCP on changes in the parameters indicative of oxidative stress in rats. The results showed increased production of reactive oxygen species, lipid peroxidation and altered serum transaminases, which might be contributing to organophosphates toxicity and tissue damage.

It is apparent that, although inhibition of cholinesterases plays a key role in the toxicology of organophosphates the inhibition of other enzyme systems and the direct effects of organophosphates on tissues are also important. Organophosphates are known to induce oxidative stress, but there is little evidence of changes in antioxidant system after intoxication with organophosphate compound (Hai *et al* 1997). Reports indicate that toxic manifestations induced by organophosphates may be associated with the enhanced production of ROS. It is also reported that different classes of pesticides induce reactive oxygen species (ROS) leading to tissue oxidative damage (Bagchi *et al* 1995). Also excessive amounts of ROS generation due to high-energy consumption coupled with inhibition of oxidative phosphorylation leads to decreased capacity of cells to maintain its energy levels (Milatovic *et al* 2006). We observed increased free radical generation in the present study predominantly in the animals given MCP alone and in combination with DDVP (Dwivedi *et al* 2011).

Reports indicate that enzyme activities associated with antioxidant defence mechanisms are altered by insecticides both in vivo and in vitro (Gultekin *et al* 2000; Oncu *et al* 2002). Both the increased production of reactive oxygen species and attenuation of the antioxidant barrier of the organism are likely to induce oxidative stress, leading to tissue damage,

tubular necrosis and cardiotoxicity in acute as well as in sub-chronic OP intoxication (Abdollahi *et al* 2004; Akhgari *et al* 2003). In the present observation also, SOD activity which is a superoxide scavenger, inhibited in MCP alone and combination groups suggesting excessive generation of reactive oxygen species followed by a parallel depletion of antioxidant enzymes. On the contrary, no changes in GSH level have been reported either in individual or in combination group suggestive of protective effect of body against the toxic effect of these toxicants.

CONCLUSION

Our findings demonstrated that monocrotophos clearly possessed the propensity to exacerbate hepatotoxicity when administered even at a relatively low dose. While the exact mechanism/s by which OPs such as monocrotophos potentiate the toxicity needs further investigation, we currently examined the possibility of oxidative stress in the augmentation of hepatotoxicity. On comparison between DDVP and MCP administered individually, MCP seems to be more potent in inducing oxidative stress. Thus, the present study provides some interesting new observations for possible co-exposure to organophosphates. On the basis of present observation it however may be suggested that co-exposure to DDVP and MCP may produce synergistic effects at some extent. The mechanism underlying their combined toxicity needs further exploration. It would also be of interest to determine the impact of OPs at further low concentrations (at levels not causing AChE inhibition) on liver degeneration.

REFERENCES

- Abdollahi M, Ranjbar A, Shadnia S, Nikfar S, Rezaie A. Pesticides and oxidative stress: a review. *Med. Sci. Monit.* 2004;10(6):RA141-7.
- Agrahari S, Pandey KC, Gopal K. Biochemical alteration induced by monocrotophos in the blood plasma of fish, *Channa punctatus* (Bloch). *Pest. Biochem. Physiol.* 2007; 88(3):268-72. [DOI: 10.1016/j.pestbp.2007.01.001]
- Akhgari M, Abdollahi M, Kebryaezadeh A, Hosseini R, Sabzevari O. Biochemical evidence for free radical-induced lipid peroxidation as a mechanism for subchronic toxicity of malathion in blood and liver of rats. *Human. Exp. Toxicol.* 2003; 22(4):205-11. [DOI: 10.1191/0960327103ht3460a]
- Aygun D. Diagnosis in an acute organophosphate poisoning: report of three interesting cases and review of the literature. *Eur. J. Emerg. Med.* 2004;11(1):55-8.
- Azmi MA, Naqvi SN, Azmi MA, Aslam M. Effect of pesticide residues on health and different enzyme levels in the blood of farm workers from Gadap (rural area) Karachi-Pakistan. *Chemosphere* 2006;64(10):1739-44. [DOI: 10.1016/j.chemosphere.2006.01.016]
- Bagchi D, Bagchi M, Hassoun EA, Stohs SJ. In vitro and in vivo generation of reactive oxygen species, DNA damage and lactate dehydrogenase leakage by selected pesticides. *Toxicology* 1995;104(1-3):129-40. [DOI: 10.1016/0300-483X(95)03156-A]
- Banerjee BD, Seth V, Ahmed RS. Pesticide-induced oxidative stress: perspectives and trends. *Rev. Environ. Health* 2001; 16(1):1-40.
- Chitra GA, Muraleedharan VR, Swaminathan T, Veeraraghavan D. Use of pesticides and its impact on health of farmers in South India. *Int. J. Occup. Environ. Health* 2006;12(3):228-33. [DOI: 10.1179/0eh.2006.12.3.228]
- Dawson AH, Eddleston M, Senarathna L, Mohamed F, Gawarammana I, Bowe SJ, Manuweera G, Buckley NA. Acute human lethal toxicity of agricultural pesticides: a prospective cohort study. *PLoS Med.* 2010;7(10): e1000357 [DOI: 10.1371/journal.pmed.1000357]

- Delescluse C, Ledirac N, Li R, Piechocki MP, Hines RN, Gidrol X, Rahmani R. Induction of cytochrome P450 1A1 gene expression, oxidative stress, and genotoxicity by carbaryl and thiabendazole in transfected human HepG2 and lymphoblastoid cells. *Biochem. Pharmacol.* 2001;61(4):399-407. [DOI: 10.1016/S0006-2952(00)00562-1]
- Dwivedi N, Bhutia YD, Kumar V, Yadav P, Kushwaha P, Swarnkar H, Flora SJ. Effects of combined exposure to dichlorvos and monocrotophos on blood and brain biochemical variables in rats. *Human Exp. Toxicol.* 2010;29(2):121-9. [DOI: 10.1177/0960327109357212]
- Dwivedi N, Flora SJ. Concomitant exposure to arsenic and organophosphates on tissue oxidative stress in rats. *Food Chem. Toxicol.* 2011;49(5):1152-9. [DOI: 10.1016/j.fct.2011.02.007]
- Eddleston M. The pathophysiology of organophosphorus pesticide self-poisoning is not so simple. *Neth. J. Med.* 2008;66(4):146-8.
- Elia AC, Galarini R, Dorr AJM, Taticchi MI. Bioaccumulation of heavy metals, organochlorine pesticides, and detoxication biochemical indexes in tissues of Ictalurus melas of Lake Trasimeno. *Bull. Environ. Contam. Toxicol.* 2006;76(1):132-9. [DOI: 10.1007/s00128-005-0899-1]
- Environmental Protection Agency US. Health and environmental effects profile for dichlorvos. 2000; 11-5.
- Ewing RD. Diminishing returns: Salmon decline and pesticides. Funded by the Oregon Pesticide Education Network, Biotech Research and Consulting, Inc., Corvallis. 1999; 55.
- Flessel P, Quintana PJE, Hooper K. Genetic toxicity of malathion: a review. *Environ. Mol. Mutagen.* 1993;22(1):7-17. [DOI: 10.1002/em.2850220104]
- Folmar LC. Effects of chemical contaminants on blood chemistry of teleost fish: a bibliography and synopsis of selected effects. *Environ. Toxicol. Chem.* 1993;12(2):337-75. [DOI: 10.1002/etc.5620120216]
- Gultekin F, Ozturk M, Akdogan M. The effect of organophosphate insecticide chlorpyrifos-ethyl on lipid peroxidation and antioxidant enzymes (in vitro). *Arch. Toxicol.* 2000;74(9):533-8. [DOI: 10.1007/s002040000167]
- Gupta PK. Pesticide exposure-Indian scene. *Toxicology* 2004;198(1-3):83-90. [DOI: 10.1016/j.tox.2004.01.021]
- Hai DQ, Varga SI, Matkovic B. Organophosphate effects on antioxidant system of carp (*Cyprinus carpio*) and catfish (*Ictalurus nebulosus*). *Comp. Biochem. Physiol. C Pharmacol. Toxicol. Endocrinol.* 1997;117(1):83-8. [DOI: 10.1016/S0742-8413(96)00234-4]
- Halliwell B, Whiteman M. Measuring reactive species and oxidative damage in vivo and in cell culture: how should you do it and what do the results mean? *Br. J. Pharmacol.* 2004;142(2):231-55. [DOI: 10.1038/sj.bjp.0705776]
- Hettwer H. Histochemical investigations on liver and kidney of the rat after intoxication with organophosphates. *Acta Histochem.* 1975;52(2):239-52.
- Hissin PJ, Hilf R. A fluorometric method for determination of oxidized and reduced glutathione in tissues. *Anal. Biochem.* 1976;74(1):214-26. [DOI: 10.1016/0003-2697(76)90326-2]
- Horrigan L, Lawrence RS, Walker P. How sustainable agriculture can address environmental and human health harms of industrial agriculture. *Environ. Health Perspect.* 2002;110(5):445-56.
- Julka D, Pal R, Gill KD. Neurotoxicity of dichlorvos: effect on antioxidant defense system in the rat central nervous system. *Exp. Mol. Pathol.* 1992;56(2):144-52. [DOI: 10.1016/0014-4800(92)90031-6]
- Kakkar P, Das B, Viswanathan PN. A modified spectrophotometric assay of superoxide dismutase. *Indian J. Biochem. Biophys.* 1984;21(2):130-2.
- Kalender S, Ogutcu A, Uzunhisarcikli M, Acikgoz F, Durak D, Ulusoy Y, Kalender Y. Diazinon-induced hepatotoxicity and protective effect of vitamin E on some biochemical indices and ultrastructural changes. *Toxicology* 2005; 211(3):197-206. [DOI: 10.1016/j.tox.2005.03.007]
- Lowry OH, Rosebrough NJ, Farr AL, Randall RJ. Protein measurement with the Folin phenol reagent. *J. Biol. Chem.* 1951;193(1):265-75.
- Milatovic D, Gupta RC, Aschner M. Anticholinesterase toxicity and oxidative stress. *ScientificWorldJournal* 2006;6:295-310. [DOI: 10.1100/tsw.2006.38]
- Mittal M, Flora SJ. Effects of individual and combined exposure to sodium arsenite and sodium fluoride on tissue oxidative stress, arsenic and fluoride levels in male mice. *Chem. Biol. Interact.* 2006;162(2):128-39. [DOI: 10.1016/j.cbi.2006.05.018]
- Naqvi SM, Hasan M. Acetylhomocysteine thiolactone protection against phosphamidon-induced alteration of regional superoxide dismutase activity in central nervous system and its correlation with altered lipid peroxidation. *Indian J. Exp. Biol.* 1992;30(9):850-2.
- Nemcsok J, Orban L, Asztalos B, Vig E. Accumulation of pesticides in the organs of carp, *Cyprinus carpio* L., at 4 degrees and 20 degrees C. *Bull. Environ. Contam. Toxicol.* 1987;39(3):370-8.
- Nishikimi M, Appaji N, Yagi K. The occurrence of superoxide anion in the reaction of reduced phenazine methosulfate and molecular oxygen. *Biochem. Biophys. Res. Commun.* 1972;46(2):849-54. [DOI: 10.1016/S0006-291X(72)80218-3]
- Ohkawa H, Ohishi N, Yagi K. Assay for lipid peroxides in animal tissues by thiobarbituric acid reaction. *Anal. Biochem.* 1979;95(2):351-8. [DOI: 10.1016/0003-2697(79)90738-3]
- Oncu M, Gultekin F, Karaoz E, Altuntas I, Delibas N. Nephrotoxicity in rats induced by chlorpyrifos-ethyl and ameliorating effects of antioxidants. *Human Exp. Toxicol.* 2002;21(4):223-30. [DOI: 10.1191/0960327102ht225oa]
- Pimentel D. Effect of pesticides on the environment. In: 10th International Congress on Plant Protection. Crydon: UK, 1983; pp. 685-91.
- Ramaneswari K, Rao LM. Influence of endosulfan and monocrotophos exposure on the activity of NADPH cytochrome c reductase (NCCR) of *Labeo rohita* (Ham). *J. Environ. Biol.* 2008; 29(2):183-5.
- Reichling JJ, Kaplan MM. Clinical use of serum enzymes in liver disease. *Dig. Dis. Sci.* 1988;33(12):1601-14.
- Reitman S, Frankel S. A colorimetric method for the determination of serum glutamic oxaloacetic and glutamic pyruvic transaminases. *Am. J. Clin. Pathol.* 1957;28(1): 56-63.
- Ryhanen R, Herranen J, Korhonen K, Penttila I, Polvilampi M, Puhakainen E. Relationship between serum lipids, lipoproteins and pseudocholinesterase during organophosphate poisoning in rabbits. *Int. J. Biochem.* 1984; 16(6):687-90. [DOI: 10.1016/0020-711X(84)90039-9]
- Saxena G, Flora SJ. Lead-induced oxidative stress and hematological alterations and their response to combined administration of calcium disodium EDTA with a thiol chelator in rats. *J. Biochem. Mol. Toxicol.* 2004;18(4): 221-33. [DOI: 10.1002/jbt.20027]

- Shugart LR, McCarthy JF, Halbrook RS. Biological markers of environmental and ecological contamination: a review. *Risk Anal.* 1992;12(3):353-60. [DOI: 10.1111/j.1539-6924.1992.tb00687.x]
- Sinha AK. Colorimetric assay of catalase. *Anal. Biochem.* 1972;47(2):389-94. [DOI: 10.1016/0003-2697(72)90132-7]
- Socci DJ, Bjugstad KB, Jones HC, Pattisapu JV, Arendash GW. Evidence that oxidative stress is associated with the pathophysiology of inherited hydrocephalus in the H-Tx rat model. *Exp. Neurol.* 1999;155(1):109-17. [DOI: 10.1006/exnr.1998.6969]
- Sultatos LG. Mammalian toxicology of organophosphorus pesticides. *J. Toxicol. Environ. Health* 1994;43(3):271-89. [DOI: 10.1080/15287399409531921]
- Taylor P. Anticholinesterase Agents. In: The Pharmacological Basis of Therapeutics, 9th edition, Hardman JG, Limbird LE (eds). McGraw Hill, New York: USA, 1996; pp. 161-76.
- Tyler CR, Jobling S, Sumpter JP. Endocrine disruption in wildlife: a critical review of the evidence. *Crit. Rev. Toxicol.* 1998;28(4):319-61.
- Waite DT, Sommerstad H, Grover R, Kerr L, Westcott ND. Pesticides in ground water, surface water and spring runoff in a small Saskatchewan watershed. *Environ. Toxicol. Chem.* 1992;11(6):741-8. [DOI: 10.1002/etc.5620110603]
- Yamano T, Morita S. Hepatotoxicity of trichlorfon and dichlorvos in isolated rat hepatocytes. *Toxicology* 1992;76(1):69-77. [DOI: 10.1016/0300-483X(92)90019-B]
- Yang W, Sun AY. Paraquat-induced free radical reaction in mouse brain microsomes. *Neurochem. Res.* 1998;23(1):47-53. [DOI: 10.1023/A:1022497319548]
